

Children's Tylenol

		Infants' Concentrated Drops 80mg/0.8ml	Children's Suspension Liquid 160mg/5ml	Children's Soft Chewable Tablets 80mg each	Junior Strength Chewable Tablets 160mg each
WEIGHT	AGE	Dropperful	Teaspoon (TSP)	Tablets	Tablets
6-11 lbs	0-3 mos	1/2= (0.4ml)			
12-17 lbs	4-11 mos	1= (0.8ml)	1/2 TSP		
18-23 lbs	12-23 mos	1-1/2	3/4 TSP		
24-35 lbs	2-3 yrs	2	1 TSP	2	
36-47 lbs	4-5 yrs		1-1/2 TSP	3	
48-59 lbs	6-8 yrs		2 TSP	4	2
60-71 lbs	9-10 yrs		2-1/2 TSP	5	2-1/2
72-95 lbs	11 yrs		3 TSP	6	3
96+ lbs	12+ yrs				4

Note: May repeat dose every 4 hours with maximum of 5 doses per 24 hour period.

Note: If possible, use weight to dose; otherwise use age.

Children's Motrin

		Infants' Motrin Concentrated Drops 50mg/1.25ml	Children's Motrin Suspension 100/ 5ml	Children's Motrin Chewable Tablets 50mg	Junior Strength Motrin Chewable Tablets 100mg
WEIGHT	AGE	Dropperful	Teaspoon (TSP)	Tablets	Tablets
6-11 lbs	0-6 mos	1/2 to 1			
12-17 lbs	6-11 mos	1			
18-23 lbs	12-23 mos	1-1/2			
24-35 lbs	2-3 yrs		1 TSP	2	
36-47 lbs	4-5 yrs		1-1/2 TSP	3	
48-59 lbs	6-8 yrs		2 TSP	4	2
60-71 lbs	9-10 yrs		2-1/2 TSP	5	2-1/2
72-95 lbs	11 yrs		3 TSP	6	3

Note: May repeat dose every 6 to 8 hours. If your physician recommends alternating Motrin and Tylenol every 2 hours, then decrease the above doses of Motrin by 1/2 to 3/4 dose. (ex: If dose is normally 1 TSP every 6-8 hrs, then 1/2 x 1 = 1/2 TSP, and 3/4 x 1 TSP = 3/4 TSP)

Benadryl Dosages

Product	Children's Benadryl Allergy Liquid	Children's Benadryl Allergy Liquid Dye-Free	Children's Benadryl-D Allergy & Sinus	Children's Benadryl Allergy Chewables	Children's Benadryl Allergy Fastmelt	Children's Benadryl Allergy & Cold Fastmelt
Symptoms	itchy, watery eyes runny nose itchy throat sneezing	itchy, watery eyes runny nose itchy throat sneezing	itchy, watery eyes runny nose itchy throat sinus pressure sneezing	itchy, watery eyes runny nose itchy throat sneezing	itchy, watery eyes runny nose itchy throat sneezing	itchy, watery eyes runny nose sinus pressure sneezing cough

DOSAGES	Children 6 to under 12 years (> 50 lbs)	1-2 tsp every 4-6 hours as needed	1-2 tsp every 4-6 hours as needed	1-2 tsp every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed	1-2 tablets every 4-6 hours as needed
	Children 2 to 5 years (35 - 50 lbs)	3/4 tsp every 6 hours as needed	3/4 tsp every 6 hours as needed	3/4 tsp every 6 hours as needed	1 tablet every 6 hours as needed	1 tablet every 6 hours as needed	1 tablet every 6 hours as needed
	Children 6 mths to 2 yrs (20 - 35 lbs)	1/2 tsp every 6 hours as needed	1/2 tsp every 6 hours as needed	1/2 tsp every 6 hours as needed	not recommended	not recommended	not recommended
	Children 3-6 months (10 - 20 lbs)	1/4 tsp every 6 hours as needed	1/4 tsp every 6 hours as needed	1/4 tsp every 6 hours as needed	not recommended	not recommended	not recommended

**** Do not exceed 4 doses in 24 hours for all products**

lbs=pounds

tsp=teaspoonful

mths=months

1 dropperful=0.8 ml

1 teaspoonful=5 ml